

Neutrino Fluxes and Resonance Physics with Neutrino Telescopes

Myron Bander*

Department of Physics, University of California, Irvine, California 92717, USA

H. R. Rubinstein**

Department of Theoretical Physics, University of Uppsala, Uppsala, Sweden

and

Department of Theoretical Physics, Hebrew University, Jerusalem, Israel

(March 1994)

Abstract

Expected atmospheric $\bar{\nu}_e$ fluxes will result in a significant number of resonantly produced low mass hadronic vector states, as the ρ , in large volume neutrino telescopes. The existence of sources of higher energy neutrinos will result in the production of higher mass states, as the D_s^* and the $(\bar{t}b)_{J=1}$. We calculate the rates of production of these states and discuss their signals. Independent of theoretical flux determinations, the detection of these states will be a tool for experimentally determining these fluxes.

PACS numbers: 96.40.Tv, 14.60.G, 98.70.Vc

Typeset using REVTeX

*Electronic address: mbander@funth.ps.uci.edu

**Electronic address: rub@vana.physto.se

Large volume high energy neutrino telescopes are coming into operation with effective areas of the order of 10^9 cm^2 and volumes greater than $2 \times 10^{13} \text{ cm}^3$ [1–3]. Planned is the NESTOR [4] detector with a potential volume of 10^{14} cm^3 . In this note we point out a method of determining or setting bounds on $\bar{\nu}_e$ fluxes at various energies corresponding to the production of the standard vector quark-antiquark resonances, namely ρ , D_s^* and possibly $(\bar{t}b)$; rates for the production of other states is significantly lower because of Cabibbo or helicity suppression. This complements the Glashow [5] mechanism for resonant W^- production.

A guaranteed source, with energies below 10^4 GeV , are atmospheric neutrinos [6]. Neutrinos in this energy range will be capable of producing the ρ meson and the D_s^* . The extrapolated, to higher energies, atmospheric neutrino fluxes give totally negligible rates for the production of higher mass resonances. However, interesting sources of neutrinos with energies higher than 10^4 GeV have been postulated; Active Galactic Nuclei (AGN) [7] are the most promising source in that TeV gamma rays have been detected from Markarian 421 [8] and there is consensus that fluxes of all neutrino flavors of comparable intensity exist [9]; as a large number of AGNs is known to exist one can estimate the expected diffuse neutrino flux [10]. Unexpected sources might be early universe relic neutrinos [11].

Some time back Glashow [5] pointed out that the W boson can be produced in a resonant way in neutrino-electron scattering. We study the resonance production of hadronic states: ρ 's, D_s^* 's with a mass of 2.1 GeV and of the $J = 1$ state of the $(\bar{t}b)$ system. For completeness we also present results for the helicity suppressed π production and the Cabibbo suppressed K^* production. For a vector meson R the rate in a volume V is

$$\text{Rate} = \frac{48\pi^3\Gamma(R \rightarrow e\nu)}{M_R m_e} \Phi\left(\frac{M_R^2}{2m_e}\right) N_e V, \quad (1)$$

where N_e is the electron density (in water $N_e = 3.4 \times 10^{23}/\text{cm}^3$) and $\Phi(E)$ is the flux, averaged over azimuthal angles, at $\bar{\nu}_e$ energy E . In the above we assume that the flux does not change rapidly over the width of the resonance.

The partial width $\Gamma[(Q\bar{q})_{J=1} \rightarrow e\nu]$, for a vector resonance made out of Q and \bar{q} quarks,

is obtained from the related electromagnetic of the $Q\bar{Q}$ system. For the latter we use the empirical relation [12]

$$\Gamma[(Q\bar{Q})_{J=1} \rightarrow e^-e^+] = 12e_Q^2 \text{keV}, \quad (2)$$

where e_Q is the charge of the Q quark. In terms of the nonrelativistic quark model this implies that the wave function at the origin is proportional to the reduced mass. Using these facts we find

$$\Gamma[(Q\bar{q})_{J=1} \rightarrow e\nu] = 192M_Q^2 M_q^2 \left(\frac{G_F}{4\pi\alpha}\right)^2 \text{keV}. \quad (3)$$

For the ρ meson conservation of isospin gives a firmer result, consistent with the one above,

$$\Gamma(\rho \rightarrow e\nu) = 12M_\rho^4 \left(\frac{G_F}{4\pi\alpha}\right)^2 \text{keV}. \quad (4)$$

Combining Eq. (1) and Eq. (3) we find for a volume of $2 \times 10^{13} \text{ cm}^3$ (the smallest of the proposed neutrino telescope volumes)

$$\text{Rate} = 8 \times 10^{11} \frac{M_Q^2 M_q^2}{M_R} \Phi \left(\frac{M_R^2}{2m_e}\right) / \text{year}, \quad (5)$$

where Φ is in units of $(\text{cm}^2 \text{ s sr GeV})^{-1}$ and all masses are in GeV. For the ρ meson the corresponding result is

$$\text{Rate} = 1.7 \times 10^{10} \Phi(580 \text{ GeV}) / \text{year}. \quad (6)$$

The results, with calculated atmospheric neutrino fluxes (ATM) [6] and theoretically estimated active galactic neutrino fluxes (AGN) [10,13] are given in Table I. For convenience we present the flux needed to obtain 10 events/year. (For the calculations we use a t quark mass of 175 GeV.)

We also present, in Table II, results for the production of two low mass states that are either helicity suppressed π or Cabibbo suppressed K^* . We note that at $E = 6.4 \times 10^6 \text{ GeV}$, the W^- production rate is 32/year.

The atmospheric neutrino flux calculations [6] are conservative and we certainly expect that the neutrino telescopes will see several ρ events per year. The ν_μ fluxes are calculated

to be an order magnitude larger. Should there be any significant neutrino mixing [14] enhancing the ν_e flux, the rates due to atmospheric neutrinos would go up significantly. As mentioned earlier the rates presented in this work are for the smallest volume telescope planned. Although the rate for production of $(\bar{t}b)_{J=1}$ due to the calculated AGN neutrino flux is well below the feasibility of any telescope, there might be totally unexpected sources. The detection of hadronic resonances will provide an experimental determination or limit on $\bar{\nu}_e$ fluxes. We have presented rates for events totally contained in the neutrino telescope volume. These events will be characterized by having no visible particle entering the volume and 600 GeV or more of hadronic energy deposited locally in the detector in thin hadronic and/or electromagnetic shower of length 6 m to 10 m. [15].

M. B. was supported in part by the National Science Foundation under Grants PHY-9208386 and INT-9224138. H. R. was supported by the Swedish Research Council and an EEC Science grant. We wish to thank Dr. S. Barwick, Dr. S. Carius and Mr. Mats Thunman for valuable discussions.

REFERENCES

- [1] S. Barwick *et al* in *Proceedings of the Workshop on High Energy Neutrino Astrophysics, Honolulu, 1992*, edited by V. J. Stenger *et. al.* (World Scientific, Singapore, 1992), p. 291.
- [2] J. G. Learned in *Proceedings of the 2nd International Workshop on Neutrino Telescopes, Venice, 1990*, edited by M. Baldo-Ceolin.
- [3] G. D. Domogatsky in *Proceedings of the 3rd International Workshop on Neutrino Telescopes, Venice, 1991*, edited by M. Baldo-Ceolin.
- [4] L. K. Resvanis in Ref. [1], p. 325.
- [5] S. L. Glashow, Phys. Rev. **118**, 316 (1960).
- [6] P. Lipari, Astroparticle Physics **1**, 195 (1993) and references therein.
- [7] A. P. Szabo and R. J. Protheroe in Ref. [1], p. 24 and references therein.
- [8] C. W. Akerlof *et al* in *Proceedings of the XXVI International Conference on High Energy Physics, Dallas, 1992*, edited by J. R. Sanford, (American Institute of Physics, New York, 1993), vol. II, p.1214.
- [9] R. J. Protheroe in *Proceedings of the TAUP 93, Conference, Gran Sasso, Italy, September 93*, to appear in Nuclear Physics B (Proceedings).
- [10] F. W. Stecker, C. Done, M. H. Salamon and P. Sommers, Phys. Rev. Lett. **66**, 2697 (1991); the flux in this reference has to be scaled down by a factor of 30, T. Stanev in Ref. [1], p. 354.
- [11] V. Berezhinsky, Neutrinos from Early Universe Relics, Gran Sasso preprint to appear in Astroparticle Physics (unpublished).
- [12] B. L. Ioffe, V. A. Khoze and L. N. Lipatov, *Hard Processes, Volume 1, Phenomenology Quark-Parton Model*, (Nort Holland, Amsterdam, 1984), p. 160.

- [13] E. Zas, F. Halzen and R. A. Vazquez, *Astroparticle Physics* **1**, 297 (1993).
- [14] D. Casper *et al.*, *Phys. Rev. Lett.* **66**, 2561 (1991); R. Becker-Szendy *et al.*, *Phys. Rev. D* **46**, 3720 (1992); *Phys. Rev. Lett.* *69*, 1010 (1992); T. Kajita in Ref. [8], p. 1187.
- [15] S. Barwick, private communication.

TABLES

TABLE I. Event Rate for Unsuppressed Vector Meson Production in $2 \times 10^{13} \text{ cm}^3$.

State	Energy(GeV)	ATM Flux	AGN Flux	Events/year	Flux for 10/yr.
ρ	5.8×10^2	6×10^{-11}		1.2	5×10^{-10}
D_s^*	4.4×10^3	10^{-13}		0.03	3×10^{-11}
$(\bar{t}b)_{J=1}$	3.2×10^7		5×10^{-21}	2×10^{-5}	2.5×10^{-15}

TABLE II. Event Rate for Suppressed Meson Production in $2 \times 10^{13} \text{ cm}^3$.

State	Energy(GeV)	ATM Flux	Events/year	Flux for 10/yr.
π	19.4	10^{-5}	0.03	3×10^{-3}
K^*	7.8×10^2	6×10^{-11}	0.06	8×10^{-9}